

Case Report

Primary central nervous system Hodgkin's lymphoma – A case report and review of literature

Botta Thrinadh, MS¹, Kavindapadi Veerasamy Karthikeyan, MCh¹, Vijayashree Raghavan, MD², Masilamani Sathyaseelan, MS¹, Vengalathur Ganesan Ramesh, MCh¹

Departments of ¹Neurosurgery and ²Pathology, Chettinad Hospital and Research Institute, Chettinad Academy of Research and Education, Kelambakkam, Chengalpattu, Tamil Nadu, India.

ABSTRACT

Hodgkin's lymphoma (HL) is primarily a disease affecting the lymph nodes. Extranodal involvement is rare. Primary central nervous system involvement in HL is extremely rare. The diagnosis is established by morphology and distinct immune-histochemical staining of the biopsies. A 76-year-old male patient presented with speech disturbance and weakness of the right upper and lower limbs. Magnetic resonance imaging of the brain showed an intraparenchymal mass lesion in the left frontal lobe with features of a high-grade tumor. Craniotomy and complete excision of the mass lesion were done. With histopathological examination and immunohistochemistry, the diagnosis of HL was made. Whole-body positron emission tomography did not reveal any other focus. The patient is being followed up with further oncological management. Only 26 cases of primary HL have been reported in the literature. Hence, this case is presented for its rarity. The previously reported cases and the recent concepts in the pathogenesis and treatment have also been reviewed.

Keywords: Central nervous system lymphoma, Hodgkin's lymphoma of central nervous system, Primary central nervous system Hodgkin's lymphoma

INTRODUCTION

Hodgkin's lymphoma (HL) commonly affects the lymph nodes. Central nervous system (CNS) lymphoma can present either as secondary CNS involvement by systemic lymphoma which is more common or rarely as primary CNS lymphoma which is restricted to the brain, leptomeninges, spinal cord, or eyes.^[1-4] CNS involvement in HL is shown to occur in <0.2–0.5% of HL patients when compared to 5–30% CNS involvement in NHL patients.^[5,6] Primary CNS lymphoma of CNS is NHL and is usually secondary to an immunocompromised state and HL has secondary CNS involvement usually in the leptomeninges. Primary CNS involvement is extremely rare and only about 26 cases have been reported so far. Due to its rarity, the exact pathogenesis, its relation to the immune status of the patient, the exact role of Epstein-Barr Virus, its activation of regulator T (T_{reg}) cells, and the role of programmed cell death ligand (PD-L) 1 in the genesis of primary CNS HL are yet to be investigated fully.

CASE REPORT

A 76-year-old male presented to the Department of Neurosurgery with behavioral change, speech disturbance,

with weakness of the right upper and lower limbs for two weeks. The patient was conscious, disoriented, and confused and responded slowly to commands. He had right hemiparesis with Grade 3 power. He had no lymphadenopathy or hepatosplenomegaly. He was not immunocompromised magnetic resonance imaging Brain showed evidence of an irregular heterogeneous contrast-enhancing lesion in the left frontal lobe with surrounding edema and massive midline shift. The features were suggestive of high-grade glioma [Figure 1]. The patient underwent craniotomy and complete excision of the left frontal lobe mass lesion. Intraoperatively, the tumor was well defined with firm margins, moderate vascularity with a good plane of cleavage from the surrounding brain and could be excised *en masse*. The squash smears showed predominantly histiocytes with emperipolesis. The background showed numerous lymphocytes, histiocytes, occasional neutrophils, and atypical large cells. Scattered glial and neuronal cells are also shown [Figure 2]. Differential diagnosis of Rosai-Dorfman disease, histiocytic neoplasms, lymphoma, and melanoma were considered. Histopathological examination (hematoxylin and eosin-stained sections) showed brain

*Corresponding author: Vengalathur Ganesan Ramesh, Department of Neurosurgery, Chettinad Hospital and Research Institute, Kelambakkam, Tamil Nadu, India. drvgramesh@hotmail.com

Received: 25 October 2024 Accepted: 09 November 2024 Epub ahead of print: 04 January 2025 Published: XXXXX DOI: 10.25259/JNRP_402_2024

This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 4.0 License, which allows others to remix, transform, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms. ©2025 Published by Scientific Scholar on behalf of Journal of Neurosciences in Rural Practice




Figure 1: MRI Brain (T1 contrast) showing irregular contrast enhancing mass lesion in the left frontal lobe with mass effect.




Figure 2: Squash smears predominantly showing histiocytes with emperipolesis, H&E, 40X. H&E: Hematoxylin and eosin.

tissue with an infiltrating tumor composed of scattered large cells showing large vesicular nuclei, prominent eosinophilic nucleoli resembling mononuclear Hodgkin's cells, Reed Sternberg (RS) cells, and popcorn cells [Figure 3]. Atypical mitosis was noted with 7–10 mitotic figures per 10 high-powered fields. Other cells include lymphocytes and histiocytes with active phagocytosis. Large areas of necrosis and broad bands of fibrosis were seen. Immunohistochemical staining CD 45 was positive in all the hematopoietic cells and negative in the large tumor cells. The large cells expressed CD 15, CD30, nuclear positivity of PAX-5, and membranous positivity of CD79a. Ki-67 was 70%. GFAP, PAN CK, and HMB 45 were negative in the tumor cells [Figure 4]. Whole-body positron emission tomography (PET) computed tomography did not reveal any extra-cranial Hodgkin's disease. Hence, this is a case of primary CNS HL in the left frontal lobe. Postoperatively, the patient made an uneventful recovery. Speech, power, right upper and lower limbs, and behavioral disturbance improved during the postoperative period. The patient was referred for further oncological treatment. The patient had improved completely at the review at two months.




Figure 3: Histopathological images of Hodgkin's lymphoma with Reed-Sternberg cells, H&E, 40X. H&E: Hematoxylin and eosin.

DISCUSSION

Classical HL is primarily affecting the lymph nodes and CNS involvement is rare with HL, accounting for 0.2–0.5% of all




Figure 4: (a) CD45 - Positive in all the hematopoietic elements, (b) CD15 - Cytoplasmic and membranous positivity of large cells, (c) CD30- Cytoplasmic and membranous positivity of large cells, (d) CD79a- Membranous positivity of large cells, (e) PAX-5- Weak nuclear positivity of large cells, (f) Ki-67: 70%, (g) HMB - 45: Negative in the tumor cells, (h) GFAP: Negative in the tumor cells, (i) Pan CK: Negative in the tumor cells, (j) EBV: Negative in the tumor cells.

cases.^[5] The incidence of HL in CNS remains exceedingly rare in the literature with only 26 cases reported so far. The first 22 cases have been summarized by Cecyn *et al.* and five more cases (including the present case) have been added since then [Table 1].^[7-11] There have been 18 males and nine females, showing a male preponderance. In 17 cases, the lesion was in the supratentorial compartment and the rest in the posterior cranial fossa. In four patients, the lesion was dura-based mimicking a meningioma. Twenty-two patients had normal immunological status and only five had some form of immune deficiency. The most common histological type has been reported as “Classical HL - not otherwise specified” (11 cases), followed by nodular sclerosis type (seven cases) and mixed cellularity (seven cases). The Epstein-Barr virus (EBV) status has been investigated only in 13 patients, and 10 patients have been reported positive for EBV. There has been no consensus on adjuvant therapy after surgery, with varying radiotherapy and/or chemotherapy regimens. The longest reported disease-free survival has been 10 years.

The primary CNS-HL is exceedingly rare and it is of interest to note that primary CNS-HL has not been included in the primary CNS (hematolymphoid tumors) lymphomas in the latest WHO Classification of tumors of the CNS (5th Edition). Immune compromised state, if present, may contribute to

the development of primary CNS HL, though in the available literature, the majority of the patients had no evidence of immune deficiency. There has been male sex preponderance in the literature. The role of EBV has been of interest of late. The presence of EBV in the RS cells seems to correlate with the increased T_{reg} cells and may contribute to the development of this tumor.^[9] However, further research is needed in this regard. This may lead to better modalities of treatment, like targeting the T_{reg} cells. About 82% of patients with classical HL have PD-L on the surface of tumor cells and EBV infection seems to increase the expression of PD-L on the RS cells.^[9] It is of interest to note that anti-programmed cell death protein antibodies (anti-PD 1) have been used in the treatment of refractory HL. In the future anti-PD 1 antibody, like sintilimab, may form the first-line drug treatment in HL due to its relative safety compared to the potentially toxic first-line drugs used at present.

The present patient was a 76-year-old non-immunocompromised man presented with features of a rapidly growing malignant neoplasm in the left frontal lobe with imaging features suggestive of a high-grade parenchymal tumor possibly glioma. The histopathology and immunohistochemistry clinched the diagnosis of HL. We could not do the EBV DNA testing and PD-L staining which could have been very useful. The whole body PET did not reveal any focus elsewhere in the body. Hence, this is a rare case of primary CNS-HL.

Table 1: Details of the cases of primary CNS-HL reported in literature.

Case	Author	Year	Age/ sex	Immunocompromised	Location of the lesion	Type	EBV	Adjuvant treatment	Outcome
1.	Sparling et al. ^[9]	1946	53/M	Unknown	Lt frontal lobe	Hodgkin's sarcoma	Not investigated	None	Died 5 days after surgery
2.	Scricker et al. ^[10]	1955	45/M	Unknown	Rt temporal lobe	Non-sarcomatous Hodgkin's granuloma	Not investigated	RT 1500 r	NED 36 months
3.	Nagashima et al. ^[11]	1980	60/M	Unknown	Falx cerebri	HL mixed cellularity type	Not investigated	None	Unknown
4.	Bender and Mayermik ^[12]	1986	34/M	Unknown	Rt frontal lobe and dura	HL nodular sclerosing type	Not investigated	RT+CT	NED 12 months
5.	Doorly et al. ^[13]	1987	51/M	No	Lt cerebellum	HL mixed cellularity type	Not investigated	RT 30 Gy	NED 12 months
6.	Ashby et al. ^[14]	1988	62/M	No	Rt frontoparietal region	HL nodular sclerosing type	Not investigated	RT 40Gy+IT CT	NED 14 months
7.	Sickler et al. ^[15]	1990	84/M	No	Rt parietooccipital region	Classical HL NOS	Not investigated	RT 35cGy	NED 8 months
8.	Clark et al. ^[16]	1992	53/F	Unknown	Rt cerebellum	HL nodular sclerosing type	Not investigated	RT 45 Gy	NED 6 months
9.	Klein et al. ^[17]	1999	54/M	No	Rt occipital lobe	HL nodular sclerosing type	Positive	RT 36 Gy+CT	NED 16 months
10.	Blagi et al. ^[18]	2000	52/M	No	Lt temporoparietal region	HL nodular sclerosing type	Negative	RT 30 Gy+5 Gy boost	NED 21 months
11.	Johnson et al. ^[19]	2000	55/F	No	Inferior aspect of tentorium cerebelli	HL nodular sclerosing type	Positive	RT 36 Gy+14 Gy boost	NED 8 months
12.	de Castro et al. ^[20]	2007	63?M	No	Lt frontoparietal region; Lt cerebellar hemisphere	Classical HL NOS	Positive	RT 40 Gy	Unknown
13.	Hwang et al. ^[21]	2007	64/F	No	Lt cerebellar hemisphere	HL mixed cellularity type	Not investigated	RT 30 Gy+6 Gy boost	NED 16 months
14.	Gerstner et al. ^[22]	2008	58/F	No	Unknown	Classical HL NOS	Positive	RT 35 Gy	NED 90.3 months
15.	Gerstner et al. ^[22]	2008	60/F	No	Cavernous sinus, Meckel's cave, Lt mesial temporal lobe	HL nodular sclerosing type	Negative	RT	NED 1 month
16.	Foo et al. ^[23]	2011	58/M	No	Lt temporal lobe	Classical HL NOS	Positive	RT 45 GY+CT	Recurrence at 14 months
17.	Kresak et al. ^[24]	2013	70/M	Known COPD, H/o sq cell ca, HIV-ve	Lt cerebellum	Classical HL NOS	Not investigated	RT	NED 10 years
18.	Kresak et al. ^[24]	2013	72/F	No	Cerebellum	Classical HL NOS	Positive	RT	NED 6 months
19.	Henkenberens et al. ^[25]	2014	47/M	Known MS 20 years on azathioprine	Cerebellum	HL mixed cellularity type	Positive	RT 20 Gy+CT (BEACOPP)	NED 9 months

(Contd...)

Table 1: (Continued).

Case	Author	Year	Age/ sex	Immunocom promised	Location of the lesion	Type	EBV	Adjuvant treatment	Outcome
20.	Mercadal <i>et al.</i> ^[26]	2015	59/M	Known ulcerative colitis on azathioprine	Rt thalamus, midbrain	Classical HL NOS	Positive	CT (methotrexate+ cytosine arabinoside)	NED 65 months
21.	Szelernej <i>et al.</i> ^[27]	2016	47/F	Known dermatomyositis on methotrexate	Lt parietal lobe	Classical HL NOS	Not investigated	RT 30 Gy+6 Gy boost	NED 110 months
22.	Cecyn <i>et al.</i> ^[7]	2017	46/M	No	Rt frontal lobe	Classical HL NOS	Negative	RT 30 Gy+CT (ABVD)	NED 87 months died due to unrelated cause Not known
23.	Godbe <i>et al.</i> ^[8]	2019	82/F	Rheumatoid arthritis, unknown skin cancer, chronic kidney disease	Lt parietooccipital region	HL mixed cellularity type	Positive	RT	NED 58 months
24.	Fu <i>et al.</i> ^[28]	2021	60/M	No	Cerebellum	HL mixed cellularity type	Positive	CT (Methotrexate, cytosine arabinoside, and sintilimab)	NED 7 years
25.	Alfaseh <i>et al.</i> ^[29]	2019	38/M	No	Cerebellum	HL mixed cellularity type	Not investigated	RT 40 Gy+10 Gy boost+CT (ABVD)	NED 4 months
26.	Wallizada <i>et al.</i> ^[30]	2024	33/F	No	Dural based left sphenoid wing lesion	Classical HL NOS	Not investigated	CT (Multiple drug combinations)	NED 2 months
27.	Present case	2024	76/M	No	Left frontal lobe	Classical HL NOS	Not investigated	CT	NED 2 months

Table adopted from: Godbe KN, Guillems EL, Benko MJ, Grider DJ, Stump MS. primary central nervous system Hodgkin lymphoma versus lymphoproliferative disorder in an asymptomatic immunocompromised patient – a case report and review of the current literature. 2019 J. Blood Lymph 9: 250. And updated. NED: No evidence of disease, NOS: Not otherwise specified, HIV: Human immunodeficiency virus, CNS: Central nervous system, HL: Hodgkin's lymphoma, CT: Computed tomography, Lt: Left, Rt: Right, RT: Radiotherapy, MS: multiple sclerosis, COPD: Chronic obstructive pulmonary disease, BEACOPP: Bleomycin sulfate, etoposide phosphate, doxorubicin hydrochloride (Adriamycin), cyclophosphamide, vincristine sulfate (Oncovin), procarbazine hydrochloride, and prednisolone, ABVD: Adriamycin, bleomycin sulfate, vinblastine sulfate, dacarbazine.

CONCLUSION

Primary CNS-HL should be kept in the differential diagnosis of intracranial malignant tumours, especially in old men, with immunocompromised state. EBV-DNA testing and PD-L staining should be performed in all cases of suspected HL to establish the diagnosis and this will also help in planning adjuvant chemotherapy.

Ethical approval

The Institutional Review Board approval is not required.

Declaration of patient consent

The authors certify that they have obtained all appropriate patient consents.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

Use of artificial intelligence (AI)-assisted technology for manuscript preparation

The authors confirm that there was no use of artificial intelligence (AI)-assisted technology for assisting in the writing or editing of the manuscript and no images were manipulated using AI.

REFERENCES

- Haldorsen IS, Espeland A, Larsson EM. Central nervous system lymphoma: Characteristic findings on traditional and advanced imaging. *Am J Neuroradiol* 2011;32:984-92.
- Mansour A, Qandeel M, Abdel-Razeq H, Abu Ali HA. MR imaging features of intracranial primary CNS lymphoma in immune competent patients. *Cancer Imaging* 2014;14:1-9.
- Chiavazza C, Pellerino A, Ferrio F, Cistaro A, Soffietti R, Rudà R. Primary CNS lymphomas: Challenges in diagnosis and monitoring. *Biomed Res Int* 2018;2018:3606970.
- Neelakantan S, Kumaran SP, Viswamitra S, Ghosal N. Myriad of MR imaging phenotypes of primary central nervous system lymphoma in a cohort of immunocompetent Indian patient population. *Indian J Radiol Imaging* 2018;28:296-304.
- Gerstner ER, Abrey LE, Schiff D, Ferreri AJ, Lister A, Montoto S, *et al.* CNS Hodgkin lymphoma. *Blood* 2008;112:1658-61.
- Torgerson S, Olteanu H, Tinguely M, Fenske TS. Central nervous system Hodgkin lymphoma: Case report and review of the literature. *J Neurooncol* 2011;102:329-34.
- Cecyn KZ, Chaves EM, Oliveira JS. Primary central nervous system involvement in classical Hodgkin's lymphoma: Case report and review of the literature. *J Blood Lymph* 2017;8:196.
- Godbe KN, Williams EL, Benko MJ, Grider DJ, Stump MS. Primary central nervous system Hodgkin lymphoma versus lymphoproliferative disorder in an asymptomatic immunocompromised patient - a case report and review of the current literature. *J Blood Lymph* 2019;9:250.
- Sparling HJ Jr, Adams RD. Primary Hodgkin's sarcoma of the brain. *Arch Pathol (Chic)* 1946;42:338-44.
- Schricker JL, Smith DE. Primary intracerebral Hodgkin's disease. *Cancer* 1955;8:629-33.
- Nagashima K, Mori S, Yoshimasu N, Takahashi K. Primary Hodgkin's disease of the falx cerebri. *Acta Neuropathol* 1980;51:161-3.
- Bender BL, Mayernik DG. Hodgkin's disease presenting with isolated craniospinal involvement. *Cancer* 1986;58:1745-8.
- Doorly TP, Farrell MA, Phillips J. Primary intracerebral Hodgkin's lymphoma. *J Neurol Neurosurg Psychiatry* 1987;50:1048-50.
- Ashby MA, Barber PC, Holmes AE, Freer CE, Collins RD. Primary intracranial Hodgkin's disease. A case report and discussion. *Am J Surg Pathol* 1988;12:294-9.
- Sickler GK, Hanson SK, Hsu SM, Papasozomenos SC. Primary intracerebral Hodgkin's disease: A case report. *Clin Neuropathol* 1990;9:143-7.
- Clark WC, Callihan T, Schwartzberg L, Fontanesi J. Primary intracranial Hodgkin's lymphoma without dural attachment: Case report. *J Neurosurg* 1992;76:692-5.
- Klein R, Müllges W, Bendszus M, Woydt M, Kreipe, Hans M, *et al.* Primary intracerebral Hodgkin's disease: Report of a case with Epstein-Barr virus association and review of the literature. *Am J Surg Pathol* 1999;23:477-81.
- Biagi J, MacKenzie RG, Lim MS, Sapp M, Berinstein N. Primary Hodgkin's disease of the CNS in an immunocompetent patient: A case study and review of the literature. *Neurooncology* 2000;2:239-43.
- Johnson MD, Kinney MC, Scheithauer BW, Briley RJ, Hamilton K, McPherson WF, *et al.* Primary intracerebral Hodgkin's disease mimicking meningioma: Case report. *Neurosurgery* 2000;47:454-7.
- De Castro AF, Júnior AS, Horta HL, Neuenschwander LC, Fonseca RP, Lima SS, *et al.* Primary intracerebral Hodgkin lymphoma. *Br J Haematol* 2007;138:562.
- Hwang CY, Song YJ, Kim DC, Choi SS, Choi YM, Kim KU. Primary cerebellar Hodgkin's lymphoma. *J Korean Neurosurg Soc* 2007;42:149-52.
- Gerstner ER, Abrey LE, Schiff D, Ferreri AJ, Lister A, Montoto S, *et al.* CNS Hodgkin lymphoma. *Blood* 2008;112:1658-61.
- Foo WC, Desjardins A, Cummings TJ. Primary intracerebral Hodgkin lymphoma with recurrence. *Clin Neuropathol* 2011;30:75-9.
- Kresak JL, Nguyen J, Wong K, Davis R. Hodgkin lymphoma of the CNS. *Neuropathology* 2013;33:658-62.
- Henkenberens C, Franzke A, Raab P, Oschlies I, Klapper W, Christiansen H. Primary EBV-positive Hodgkin's lymphoma of the CNS under azathioprine treatment. *Strahlenther Onkol* 2014;190:847-52.
- Mercadal S, Climent F, García N, González-Barca E. Long term remission of a primary intracerebral Hodgkin lymphoma in a patient previously treated with azathioprine. *J Integr Oncol* 2015;4:144.

27. Szelemej PA, Bigder MG, Krcek J. Treatment and long-term follow-up of primary CNS classical Hodgkin's lymphoma - A case report and review of the literature. *Interdisciplin Neurosurg* 2017;9:30-3.
28. Fu H, Shi S, Chen L, Xu B, Huang W, Chen Y, *et al.* Primary central nervous system Hodgkin's lymphoma: A case report. *J Int Med Res* 2021;49:300060521999533.
29. Alfaseh A, Rajeh MN, Hamed G. Primary central nervous system Hodgkin Lymphoma: A case discussion and a hypothesis on the etiology. *Avicenna J Med* 2019;9:28-31.
30. Walizada K, Arshad MM, Khan M, Al Sulaiti GS. Classic central nervous system Hodgkin lymphoma masquerading as left sphenoid wing meningioma-A case report. *Int J Surg Case Rep* 2024;116:109439

How to cite this article: Thrinadh B, Karthikeyan KV, Raghavan V, Sathyaseelan M, Ramesh VG. Primary central nervous system Hodgkin's lymphoma – A case report and review of literature. *J Neurosci Rural Pract.* doi: 10.25259/JNRP_402_2024